

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>C12Q 1/68</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 99/54501</b> <b>(43) International Publication Date:</b> 28 October 1999 (28.10.99)
<b>(21) International Application Number:</b> PCT/IE98/00030 <b>(22) International Filing Date:</b> 22 April 1998 (22.04.98) <b>(71) Applicants (for all designated States except US):</b> ENTERPRISE IRELAND trading as BIORESEARCH IRELAND (IE/IE); Glasnevin, Dublin 9 (IE). UNIVERSITY COLLEGE CORK (IE/IE); College Road, Cork (IE). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> McCARTHY, Thomas, Valentine (IE/IE); Vista Villa, Montenotte, Cork (IE). VAUGHAN, Patrick, Martin (IE/IE); 175 West Avenue, Parkgate, Frankfield, Cork (IE). <b>(74) Agent:</b> ANNE RYAN & CO.; 60 Northumberland Road, Ballsbridge, Dublin 4 (IE).		<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> A METHOD FOR THE CHARACTERISATION OF NUCLEIC ACID MOLECULES INVOLVING GENERATION OF EXTENDIBLE UPSTREAM DNA FRAGMENTS RESULTING FROM THE CLEAVAGE OF NUCLEIC ACID AT AN ABASIC SITE		
<b>(57) Abstract</b> <p>A method for characterising nucleic acid molecules comprises the steps of: i) introducing a modified base which is a substrate for a DNA glycosylase into a DNA molecule; ii) excising the modified base by means of said DNA glycosylase so as to generate an abasic site; iii) cleaving the DNA at the abasic site so as to generate an upstream DNA fragment that can be extended; and iv) incubating the extendible upstream fragment in the presence of an enzyme, for example a polymerase or a ligase, allowing for extension thereof and a template nucleic acid and analysing the resultant fragment(s). The invention provides a novel, versatile and simple method using the above-mentioned extendible upstream DNA fragments which allows characterisation of nucleic acids and which has advantages over existing methods. One of the most important uses (but not the only use) of the method according to the invention is to scan or check a fragment of DNA (target nucleic acid) for the presence or absence of a mutation.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LJ	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

### Description

A method for the characterisation of nucleic acid molecules involving generation of extendible upstream DNA fragments resulting from the cleavage of nucleic acid at an abasic site

#### 5    Technical Field

The present invention relates to a method for characterising nucleic acid molecules which involves generating extendible upstream DNA fragments which result from the cleavage of nucleic acid at an abasic site.

#### Background Art

- 10           Characterisation of target nucleic acids is highly important for several reasons relating to confirmation of the presence or absence of a gene in a sample, confirmation of part or all of a nucleic acid sequence, and screening for the presence of known and unknown disease causing mutations leading to inherited disease and natural variations in DNA.
- 15           Although there are many known methods for characterising nucleic acids and for the detection of unknown sequence changes, the increasing amount of new genetic information being generated makes it important to develop new, better and faster methods for characterisation of nucleic acids.

- 20           WO 97/03210 discloses the use of a DNA-glycosylase enzyme, which recognises a modified base, for the direct detection of known and unknown mutations in a target nucleic acid sample. The method typically involves amplifying a target nucleic acid sample using a combination of normal DNA precursor nucleotides and one or more modified precursor nucleotide(s) where the modified precursor nucleotide replaces one of the
- 25           normal precursor nucleotides which is a substrate for a DNA glycosylase. Following excision of the modified base by the glycosylase, the resulting abasic site is cleaved and the products of the cleavage are analysed. This

method allows detection of mutations at candidate loci. However, the method of WO 97/03210 has certain limitations. For example, with this method it is not possible to detect sequence differences between nucleic acid molecules without detecting sequence similarities and thus multiple samples cannot be combined for simultaneous analysis.

#### Disclosure of Invention

The invention provides a method for characterising nucleic acid molecules, which comprises the steps of:

- i) introducing a modified base which is a substrate for a DNA glycosylase into a DNA molecule;
- ii) excising the modified base by means of said DNA glycosylase so as to generate an abasic site;
- iii) cleaving the DNA at the abasic site so as to generate an upstream DNA fragment that can be extended; and
- iv) incubating the extendible upstream fragment in the presence of an enzyme allowing for extension thereof and a template nucleic acid and analysing the resultant fragment(s).

The invention provides a novel, versatile and simple method using the above-mentioned extendible upstream DNA fragments which allows characterisation of nucleic acids and which has advantages over existing methods as indicated in the following description.

One of the most important uses (but not the only use) of the method according to the invention is to scan or check a fragment of DNA (target nucleic acid) for the presence or absence of a mutation. The method essentially consists of i) the generation of the extendible upstream DNA

fragments and ii) the subsequent use of these fragments in analysing a piece of DNA (e.g. detecting a mutation).

Preferably, the modified base used is uracil, hypoxanthine or 8-OH guanine.

- 5            Preferably, the modified bases are derived from modified precursor nucleotides which when incorporated into DNA generate said modified bases.

10           Thus, the preferred modified precursor nucleotides are dUTP, dITP and 8-OH dGTP which when incorporated into DNA generate the glycosylase substrate bases uracil, hypoxanthine and 8-OH guanine, respectively. Each of the modified precursor nucleotides is a base sugar phosphate comprising said base and a sugar phosphate moiety. Uracil in DNA is recognised specifically by uracil DNA-glycosylase (UDG) and released from DNA. UDG also recognises other uracil related bases when  
15           present in DNA. Hypoxanthine is recognised specifically by alkylpurine DNA-glycosylases (ADG) and released from DNA. This enzyme also recognises and releases N3 methyladenine, N3 methylguanine, O<sup>2</sup> methylcytosine and O<sup>2</sup> methylthymine when present in DNA. 8-OH  
20           guanine is recognised specifically by formamido-pyrimidine DNA-glycosylase (FPG) and released from DNA. This enzyme also recognises and releases ring opened purines when present in DNA. Thymine DNA glycosylase recognises and releases uracil and thymine positioned opposite guanine bases in DNA.

- 25           Modified precursor nucleotide(s) as used herein refers to a modified nucleotide or nucleotides that can be incorporated into a nucleic acid so that a modified base or bases is generated which is/are recognised and can be excised by a DNA glycosylase enzyme.

Following the introduction of the modified base, the DNA product is treated with a suitable DNA glycosylase enzyme which recognises and releases the glycosylase substrate base present in the target sample and consequently generates an apurinic or apyrimidinic (AP) site, depending on the nature of the modified base(s). AP site is the term given to a site in DNA where the base moiety of a nucleotide has been lost or removed, leaving behind a deoxyribophosphate with the DNA phosphodiester backbone still intact. AP is the abbreviation for either apurinic and/or apyrimidinic depending on whether a purine or pyrimidine base had been attached to the ribose ring. An AP site is also referred to as an abasic site, being the general term for apurinic and apyrimidinic site.

Release of the glycosylase substrate base from the nucleic acid sample results for example in an apyrimidinic site in the case of uracil and an apurinic site in the case of hypoxanthine and 8-OH dG. Collectively, such sites are referred to as abasic sites.

Essentially, cleaved fragments must have hydroxyl groups at the 3' termini and the DNA immediately downstream of said 3' termini must not be blocked in a way that prevents extension of the fragment from said 3' termini. Cleaved fragments are generated that have hydroxyl at the 3' termini and downstream blocking groups are removed that prevent extension of the fragment from the said 3' termini, while using as template, that DNA from which the extendible fragment was derived.

The DNA can be cleaved in a number of ways at the abasic site so as to generate said upstream DNA fragment as hereinafter described in greater detail.

For example, the phosphate linkages at the abasic sites can be cleaved by a treatment selected from treatment with a basic solution or other chemical treatment, heat treatment and/or treatment with an enzyme.

According to one embodiment of the invention, the upstream fragment is generated by cleaving the DNA at the 5' side of the abasic site, such that the 3' terminus of the upstream fragment bears a hydroxyl group.

- 5        The terms extendible fragment and upstream DNA fragment are used interchangeably herein.

As the 3'-OH termini are thereby generated no further processing of the upstream fragments is required prior to step iv).

- 10        Treatment with basic solutions (alkali) at high temperature or with a chemical such as piperidine, or with an enzyme which cuts specifically at abasic sites, such as *E. coli* endonuclease IV results in cleavage of the abasic site on the 5' side.

Suitably in this embodiment, the cleavage is achieved with a 5' AP endonuclease.

- 15        According to an alternative embodiment the upstream fragment is generated by cleaving at the 5' side of the abasic site so as to leave a phosphate group at the 3' terminus of the upstream fragment and removing the phosphate group so that the upstream fragment bears a hydroxyl group at the 3' terminus.

- 20        Alternatively, cleavage with alkali or the abasic endonuclease activity of FPG is used, followed by removal of the 3' phosphate. For example 3' phosphate groups can be removed by enzymatic means using enzymes with 3' phosphate activity such as T4 polynucleotide kinase.

- 25        According to a still further embodiment, the upstream fragment is generated by cleaving at the 3' side of the abasic site so as to generate a deoxyribose group phosphate at the 3' terminus of the upstream fragment



and subsequently removing the deoxyribose group to leave a hydroxyl group at the 3' terminus.

This can be achieved by using an enzyme with 3' deoxyribo-phosphodiesterase activity, or using FPG followed by a 3' phosphatase.

- 5        In an alternative embodiment 5' deoxyribose moieties downstream of the 3' terminus of the upstream fragment are removed so that the upstream fragment can be extended on the template.

Preferably the 5' deoxyribose moieties are removed by a 5' deoxyribophosphodiesterase.

- 10       Treatment with high temperature alone or a 3' AP endonuclease results in cleavage of the abasic site to completion on the 3' side.

- 15       Glycosylase mediated cleavage cuts the extended primer at an abasic site subsequent to release of the modified base by a DNA glycosylase yielding 3' termini with 3'-OH or 3' phosphate groups or deoxyribophosphate groups. Except in cases where 3'-OH termini are generated, all other termini require further processing prior to extension of the upstream fragment.

- 20       Glycosylase mediated cleavage in the method according to the invention refers to both 5' and 3' cleavage, including whatever subsequent treatment that is necessary to generate a 3' OH group at the 3' terminus of the upstream fragment.

Preferably, the modified base is introduced by enzymatic amplification of the DNA.

Preferably, the DNA (herein also referred to as target nucleic acid) is amplified using normal DNA precursor nucleotides and at least one modified precursor nucleotide.

5 Precursor nucleotides in the case of a DNA amplification process refer to the deoxyribonucleotides dATP, dCTP, dGTP and dTTP herein referred to as "normal" DNA precursor nucleotides.

10 The term "amplifying" as used herein refers to any *in vitro* process for increasing the number of copies of a nucleotide sequence or sequences. Amplification of a target nucleic acid molecule results in the incorporation of precursor nucleotides into the DNA being amplified. Typically, amplification of a target sample is carried out using appropriate primers in the polymerase chain reaction (PCR). During the amplification, primers anneal to the target nucleic acid and are extended using a DNA polymerase in a 5' to 3' direction on the target nucleic acid which acts as a template for synthesis of new DNA. The use of flanking primers, which are referred to herein as initial primers, which anneal to the upper and lower strands of a DNA molecule permit exponential amplification of the DNA segment delimited by the upper and lower primers.

20 The amplification will typically involve amplifying a target nucleic acid using a combination of normal DNA precursor nucleotides and one or more modified precursor nucleotide(s) where the modified precursor nucleotide replaces all or a proportion of one of the normal precursor nucleotides. Amplification of a nucleic acid using normal DNA precursor nucleotides results in the incorporation of the four normal bases G,A,T, or C into DNA. Amplification of a nucleic acid using a modified precursor nucleotide in place of one of the normal precursor nucleotides results in the incorporation of a glycosylase substrate base into DNA in place of one of the four normal bases G,A,T, or C.

The target nucleic acid sample will typically be DNA. However, RNA may also be used following conversion to DNA by reverse transcription.

5           When a modified precursor nucleotide replaces a proportion of one of the normal precursor nucleotides, the ratio of the modified precursor nucleotide to the normal precursor nucleotide that it is replacing is such that an optimum of one modified precursor nucleotide is incorporated *per* strand of amplified DNA. This allows subsequent cleavage of the  
10           amplified DNA strand into two fragments following contact with a DNA glycosylase and an abasic site cleavage agent as herein described. This method of cleavage is herein referred to as glycosylase mediated cleavage. A higher ratio of the modified precursor nucleotide to normal precursor nucleotide is used to generate more than one cleavage site *per* amplified  
15           DNA strand. The incorporation of a modified precursor nucleotide into the amplified product generates one or more modified bases at one or more positions recognised by a DNA glycosylase enzyme in the amplified product.

20           Replacement of all of a normal precursor nucleotide with a modified precursor nucleotide in the amplification reaction if used in step i) permits glycosylase mediated cleavage of a primer extended in an amplification reaction at the first position 3' of the extended primer where a normal base is replaced by a modified base. Thus, if the template  
25           sequence immediately 3' of a location where the primer hybridises is CTAG and the modified nucleotide precursor is dUTP replacing dTTP, then the modified base uracil (U) will be incorporated opposite A on the template strand. Thus, in this situation, the primer will have been extended by two nucleotides at the 3' end (primer - GA 3') following amplification  
30           and glycosylase mediated cleavage. These extended primers generated following the initial extension and glycosylase mediated cleavage are referred to herein *inter alia* as extendible fragments, as indicated above,

whereas the primers prior to extension are referred to as the initial primers herein.

5       The extended 3' terminal sequence of the extendible fragment is enzymatically synthesised and is directly related to the nucleic acid being characterised as the nucleic acid acts as the template for its synthesis. Thus the 3' end of the extendible fragment is complementary to the nucleic acid. Accordingly, determination of the nature of the 3' end of the extendible fragment by any means allows characterisation of the nucleic acid from  
10       which it was derived. If an initial primer is placed adjacent to a locus where a DNA sequence variation such as a polymorphism or a mutation occurs so that the first modified base incorporated into the extended primer is at the mutation locus, then the initial primer will be extended to a different length depending on whether or not a mutation is present at the  
15       mutation locus following glycosylase mediated cleavage. The extendible fragments are subsequently treated (if necessary) so that they can be used as primers for a subsequent extension reaction. Because the sequence at the 3' termini of the extendible fragments differ depending on whether a  
20       mutation is present or absent at the mutation locus, analysis of the ability of an extendible fragment to function in a subsequent extension reaction using a template nucleic acid permits the determination of whether a mutation is present or absent at the mutation locus. Any naturally  
25       occurring or enzymatically or chemically synthesised template which fully or partially hybridises to the extendible fragment can be designed and/or selected as a template nucleic acid allowing the ability of the extendible fragment to function as a primer to be determined.

30       When a proportion of a normal precursor nucleotide is replaced with a modified precursor nucleotide in the amplification reaction, glycosylase mediated cleavage of the primer extended in an amplification reaction will yield a population of extendible fragments of various lengths since different molecules will be cleaved at different points depending on where the modified precursor nucleotide is incorporated. The length of

each fragment is determined by the position of incorporation of the modified precursor nucleotide during extension from the 3' end of the initial primer.

Amplification of a target nucleic acid using the precursor  
5 nucleotides dATP, dCTP, dGTP, dTTP and a low amount of the modified  
precursor nucleotide dUTP results in an amplified DNA where thymine is  
replaced randomly by uracil. The uracil is incorporated in the newly  
synthesised DNA strand at positions complementary to adenine residues in  
10 the template DNA strand during the amplification process. Amplification  
of a target nucleic acid using the precursor nucleotides dATP, dCTP,  
dGTP, dTTP and a low amount of the modified precursor nucleotide dITP  
results in an amplified DNA where guanine is preferentially replaced  
randomly by hypoxanthine. The hypoxanthine is incorporated in the newly  
15 synthesised DNA strand at positions complementary to cytosine residues in  
the template DNA strand during the amplification process when the other  
precursor nucleotides are not limiting. Amplification of a target nucleic  
acid using the precursor nucleotides dATP, dCTP, dGTP, dTTP and a low  
amount of the modified precursor nucleotide 8-OH dGTP results in an  
20 amplified DNA where guanine is preferentially replaced randomly by 8-  
OH guanine. The 8-OH guanine is incorporated in the newly synthesised  
DNA strand at positions complementary to cytosine residues in the  
template DNA strand during the amplification process when the other  
precursor nucleotides are not limiting.

The amplified DNA strands can be separated for separate analysis of  
25 the respective strands. In addition, the separated strands can be  
immobilised, which can be achieved by several means. A common method  
for immobilisation and/or separation of DNA strands is by the use of the  
biotin streptavidin interaction, where normally, the DNA contains the  
biotin label and the streptavidin is attached to a solid support. However,  
30 the method according to the invention in its various steps is amenable to  
immobilisation formats that allows immobilisation of the upstream

fragment, the strand bearing the upstream fragment, the strand  
complementary to the strand bearing the upstream fragment, the template  
nucleic acid, the target nucleic acid and the products generated from  
glycosylase mediated cleavage of amplified or extended nucleic acids  
5 bearing modified bases.

The modified base can be introduced by chemical modification of a  
nucleic acid, rather than by an amplification technique such as PCR.

Several methods exist where the treatment of DNA with specific  
10 chemicals modifies existing bases so that they are recognised by specific  
DNA glycosylase enzymes. For example, treatment of DNA with  
alkylating agents such as methylnitrosourea generates several alkylated  
bases including N3-methyladenine and N3-methylguanine that are  
recognised and excised by alkyl purine DNA-glycosylase. Treatment of  
15 DNA with sodium bisulfite causes deamination of cytosine residues in  
DNA to form uracil residues which are recognised and excised by uracil  
DNA-glycosylase. Treatment of DNA with ferrous sulphate and EDTA  
causes oxidation of guanine residues in DNA to form 8-OH guanine  
residues in DNA which are recognised and excised by formamido-  
20 pyrimidine DNA glycosylase.

Thus, bases present in the nucleic acid or indeed, the extendible  
upstream fragment generated in step iii) can be converted into modified  
bases by chemical means. A proportion or all of the cytosine residues can  
25 be readily converted to an uracil using sodium bisulfite thereby rendering  
the amplified sample susceptible to uracil DNA-glycosylase cleavage at the  
sites of cytosine conversion. If the upper or lower primer is synthesised so  
that it contains 5-methylcytosine rather than cytosine, in such a case the  
primer will be resistant to uracil DNA-glycosylase mediated cleavage  
30 since deamination of 5-methylcytosine occurs at a reduced rate by  
comparison with cytosine and generates a thymine rather than an uracil  
residue.

Prior to treatment with a suitable DNA glycosylase, double stranded DNA may be treated with exonuclease I. This treatment serves to digest the unused primers and any non-specific single stranded DNA amplification products thus improving the signal to noise ratio.

5

In the case where the modified precursor nucleotide is dUTP, the modified base uracil will be generated at thymine positions in the amplified target nucleic acid sample. Addition of uracil DNA-glycosylase to the sample releases the uracil from the sample. In the case where the modified precursor nucleotide is dITP, the modified base hypoxanthine will be generated at guanine positions in the amplified target nucleic acid sample. Addition of alkylpurine DNA-glycosylase to the sample releases the hypoxanthine from the sample. In the case where the modified precursor nucleotide is 8-OH dGTP, the modified base 8-OH guanine will be generated at guanine positions in the amplified target nucleic acid sample. Addition of formamido-pyrimidine DNA-glycosylase to the sample releases the 8-OH guanine from the sample.

10

15

Suitably a primer or one or more nucleotide(s) involved in the enzymatic amplification is labelled.

20

The initial primer used may be suitably labelled. Labelling of the primers can be performed by a variety of means including addition of a radioactive, fluorescent, or detectable ligand to the primer during or post primer synthesis.

25

In one embodiment of the invention, the enzyme used in step iv) is a polymerase which can be incubated with the extendible upstream fragment in the presence of one or more nucleotide(s).

Also in this embodiment suitably one or more of the nucleotide(s) of step iv) is a dideoxy nucleotide.

Also one or more of the nucleotides of step iv) can be labelled.

5 Various nucleic acid polymerases can be used to extend the 3' terminus of the extendible fragment on a template nucleic acid. Many polymerases are described in the literature that extend 3' termini of primers on a template DNA. For example, DNA polymerases isolated from phage and mesophilic and thermophilic bacteria can be used.

10 Several DNA polymerases, including T7 DNA polymerase, incorporate dideoxy terminator nucleotides as well as normal precursor nucleotides during extension of primers. Thermophilic DNA polymerases are used routinely in amplification of nucleic acids through repeated cyclic extension of primers. The upstream DNA fragments generated in step iii) function as primers for all nucleic acid polymerases capable of extending standard nucleic acid primers.

15 The use of a labelled precursor nucleotide or dideoxy terminator nucleotide in any of the extension reactions facilitates detection of the extended extendible fragment. Direct DNA staining methods such as silver or ethidium bromide staining facilitate detection of all extension products after size separation based on electrophoretic mobility.

20 The ability of the extendible fragments to function in a subsequent extension reaction using a template nucleic acid and normal or dideoxy terminator nucleotides (a nucleotide that prohibits further extension of a primer on a template once incorporated) generates a ladder of fragments allowing determination of the location of the total number of positions of the modified precursor nucleotide in one or both strands of the amplified target nucleic acid. The presence of a sequence variation or mutation results in the appearance or disappearance of a cleavage fragment as judged by comparison with the known DNA sequence of the amplified molecule. Size analysis of the fragments allows the precise location and sequence of a mutation in the target nucleic acid sample to be determined.



Therefore, if a sequence variation occurs such that an additional site of modified precursor nucleotide incorporation is generated, an additional cleavage fragment will be observed upon analysis of the ladder of cleavage products. If a sequence variation occurs such that a site of modified precursor nucleotide incorporation is lost, the corresponding cleavage fragment will not be observed upon analysis of the ladder of cleavage products.

The template of choice in this case will be the originally intact or glycosylase cleaved amplified nucleic acid. In cases where glycosylase cleaved amplified nucleic acid is used it can be processed to remove residual moieties downstream of the extendible fragment which prohibit extension of the extendible fragment by a nucleic acid polymerase. Specifically the template DNA can be treated so that a residual 5' deoxyribose moiety is removed. This is achieved by incubation of the template DNA with a 5' deoxyribosephosphodiesterase such as *E. coli* RecJ endonuclease or formamidopyrimidine DNA glycosylase. Other naturally occurring or enzymatically or chemically synthesised template nucleic acids which fully or partially hybridise to the extendible fragment can also be designed and/or selected as a template nucleic acid to determine the ability of the extendible fragment to function as a primer.

The ability of the extendible fragments to function in a subsequent extension reaction using a template nucleic acid and a combination of non-labelled and labelled normal or dideoxy terminator nucleotides permits detection of sequence variations and mutations. Extension of the extendible fragments on the template nucleic acid from which they were derived and which is heterozygous for the sequence variation, using a labelled dideoxy terminator nucleotide having base pairing properties different to those of the modified precursor nucleotide with a non-labelled dideoxy terminator nucleotide having the same base pairing properties as the modified precursor nucleotide allows detection of the variant or mutant loci alone whereas non variant loci are not detected. This aspect of the invention is

particularly advantageous as detection of sequence variations alone permits very high throughput mutation scanning and detection, and allows fingerprinting of nucleic acids based on their sequence variations.

5 It will be appreciated that amplification of any target DNA, which is heterozygous for a mutation or polymorphism generates four distinct duplex DNAs, i.e. (taking as an example a G to A mutation at position X in a DNA sequence), one quarter will be homoduplex with a GC base pair at position X, one quarter will be homoduplex with an AT base pair at position X, one quarter will be heteroduplex with a GT base pair at  
10 position X, and one quarter will be heteroduplex with a AC base pair at position X. Similarly, heteroduplex DNA can be generated readily by denaturing and reannealing two homoduplex DNAs bearing a sequence difference(s).

15 Thus, the nature of the sequence of the 3'end of the extendible fragments can be determined by their ability to function as primers in a subsequent extension reaction using a template nucleic acid. Essentially, such determination is based on the ability of the 3'end of the extendible fragment to hybridise to a selected template under selected conditions. Following partial or complete hybridisation the extendible fragment may  
20 be extended using a nucleic acid polymerase and nucleic acid precursors or selected combinations of same as herein described. It will be appreciated that multiple possibilities exist for the selection of template molecules. Nonetheless, the extension of the extendible fragment is a measure of its hybridisation to or lack of hybridisation to a selected  
25 template molecule and thus the determination of the nature of the sequence of the 3'end of the extendible fragments is made on this basis, since this 3' sequence is indicative of the sequence of the original target nucleic acid.

30 Typically, the template molecule is selected so that it bears partial or full sequence complementarity to the upstream fragment. The upstream fragment may be extended one or more nucleotides on the template

molecule using a nucleic acid polymerase and nucleic acid precursors or a combination of same or dideoxy terminator nucleotides or a combination of normal nucleic acid precursors and dideoxy terminator nucleotides.

- 5       The extension of step iv) can be achieved by means of an amplification reaction using said extendible DNA fragment.

Alternatively, the extension of step iv) is achieved by means of an amplification reaction including a primer in addition to using said extendible DNA fragment.

- 10       Repeated extension of an upstream fragment on a template nucleic acid in combination with a second flanking primer which can be extended on the copy of the template permits amplification of the template nucleic acid. Such amplified products can be readily detected by standard nucleic acid staining methods such as ethidium bromide after resolution by electrophoresis.

- 15       Alternatively, the template molecule may be selected so that it can be extended using the upstream fragment as a template and so that the extension is based on hybridisation to the 3' end of the upstream fragment.

- 20       Also, the upstream fragment may be analysed based on its ability to function in a 5' nuclease assay. During extension of the upstream fragment by a polymerase with 5' to 3' nuclease activity, the 5' to 3' nuclease activity degrades a downstream reporter molecule annealed to the same template strand as said upstream fragment.

- 25       A further possibility is for the upstream fragment to be extended on a synthetic template which contains reporter and quencher labels, then cleavage of the resulting double stranded DNA will release the reporter from the quencher and a signal will be detected. Typically such cleavage

will be carried out by an enzyme which recognises the double stranded DNA molecule. Typically such an enzyme will be a restriction enzyme.

Furthermore, the 3' terminal sequence of the upstream fragment may be determined on the basis of its hybridisation to other nucleic acid molecules.

A further possibility is that the products extended or amplified using upstream fragments may be detected on the basis of their filtration and/or precipitation properties.

When analysing extension and incorporation of nucleotides in step iv) where the upstream fragments are used in an extension reaction, it is important to verify that any extension observed is specifically due to extension of the upstream fragments and not due to extension of the initial primers which were unused during the initial amplification, if such is used in step i), or extension of possible non-specific upstream fragments which may be generated by the non-specific breakage or damage of the DNA during previous steps of the procedure. To overcome this 'noise', the DNA, prior to glycosylase mediated treatment, can be treated with a single strand specific DNA nuclease, for example, Exonuclease I, which will degrade the unused primers and non-specific single stranded DNA and can be subsequently heat inactivated. The DNA can be treated with a 3' AP endonuclease/lyase which will cut the DNA and primers at any preexisting AP sites, generated specifically or generated through damage of the DNA. The 3' AP endonuclease/lyase is subsequently removed from the reaction. Because the endonuclease/lyase cuts at the 3' side, the resulting contaminating upstream fragments are not extendible and will not interfere with the extension of the subsequently generated upstream fragments. In addition to these treatments, a control reaction to check for non-specific extensions can be carried out. Thus, in step iii) of the method, the AP sites can be cut with a 3' AP endonuclease/lyase, thereby generating non-extendible upstream fragments. If however, extension and

incorporation of nucleotides is observed in the subsequent step iv) then one can measure or determine the level of non-specific extension obtained during the procedure.

5           Similarly it is important to ensure that the incorporation of nucleotides, labelled or unlabelled, in step iv) is due to the incorporation of those nucleotides supplied during step iv) and not those from any previous step. This is especially important when the reaction involves incorporation of dideoxynucleotides. Prior to or subsequent to cleaving  
10 the DNA at abasic sites and generating upstream fragments, the reaction can then be treated with a phosphatase which digests all unincorporated nucleotides present in the reaction, e.g. shrimp alkaline phosphatase, which can subsequently be heat inactivated.

15           In a further embodiment of the invention, the enzyme used in step iv) is a ligase which can be incubated with the extendible upstream fragment in the presence of a reporter oligonucleotide.

The reporter oligonucleotide may be partially degenerate.

20           The method according to the invention can be used *inter alia* to detect a known or unknown mutation and to detect differences and similarities in genomes. These aspects of the invention are illustrated further below.

25           The method according to the invention provides in one aspect a means of generating random primers in a simple easy manner. Essentially, introduction of a modified base into an amplified DNA product followed by glycosylase mediated cleavage and subsequent treatment of the cleavage products so that they can be extended by a nucleic acid polymerase provides a rapid means of generating random primers. Subsequent use of such primers, i.e., the extendible upstream fragments, in random amplification of target nucleic acids allows amplification of discrete

DNA molecules from the target nucleic acids thus permitting characterisation of the nucleic acid based on similarities and differences of the amplified products. Since these primers are essentially derived from said target nucleic acid, specifically their 3' ends, they are better primers for the subsequent random amplification analysis of said target nucleic acid and the amplification is more specific. Many discrete DNA products are generated during random amplification of nucleic acids. A discrete DNA product can be separated from other products on the basis of size. The method according to the invention permits the generation of primers from all or part of the separated product. Use of an initial primer in the random amplification of a nucleic acid that permits immobilisation of the separated product allows the isolation of the upper and lower primers extended to the first point of glycosylase mediated cleavage. The 3' end of such upstream fragments are derived from the target nucleic acid and thus permit more specific amplification of the target nucleic acid or related nucleic acids. The 3' end of such upstream fragments may be short or long. By short 3' ends herein is meant one to three nucleotides whereas by long 3' ends herein is meant greater than three nucleotides. Longer 3' ends on such upstream fragments are more desirable as they allow highly specific amplification of a target nucleic acid sequence. Upstream fragments generated with longer 3' ends may be selected by sizing methods. Alternatively, the initial amplification primers can be designed so that they promote binding of a protein that protects a section of the region 3' of the initial extended primer. Thus such a region is refractory to glycosylase mediated cleavage due to protection by the protein and inclusion of such a primer design and protein allows generation of upstream fragments with longer 3' ends.

This embodiment of the invention provides a rapid and simple method for generation of random and specific primers for nucleic acid amplification without prior knowledge of the nucleic acid sequence.

It will be appreciated that the random amplification of nucleic acid using arbitrarily chosen primers is a method already known for detecting similarities and differences between genomes. Such random amplification is based on the annealing of arbitrarily chosen primers to target samples followed by multiple rounds of enzymatic amplification whereby the primers are extended using the selected genomic DNA or cDNA as template. Using such primers in such a method results in the amplification of discrete DNA molecules. Analysis of such molecules allows the investigation of similarities and differences between different samples. Typically many different random primers are chosen for investigation of a genome or cDNA and such primers are synthesised chemically and are designed in a random fashion with the assumption that they will hybridise to the target nucleic acid in the amplification process. The method according to the invention allows the easy and rapid generation of primers from a target nucleic acid which can subsequently be used for random amplification of the same or different target nucleic acid.

In addition to extension by a polymerase reaction, as indicated above, the upstream fragments in step iv) can also be extended by ligation of another single stranded DNA molecule which results in extended upstream fragments greater in size than the initial upstream fragments generated by glycosylase mediated cleavage. The DNA molecule to which the upstream fragment is ligated is termed the reporter oligonucleotide herein and this can vary in length. Ligation of the upstream fragments to the reporter oligonucleotide is dependent on both DNA molecules (the upstream fragment and reporter oligonucleotide) annealing to a template molecule at adjacent sites so that the termini of the upstream fragment and reporter oligonucleotide are juxtaposed. This means that the 3' terminal base of the upstream fragment is juxtaposed to the 5' terminal base of the reporter oligonucleotide. The reporter oligonucleotide is typically a synthetic oligonucleotide, but can also be any other type of DNA molecule or RNA molecule.

The template molecule is selected so that it bears partial or full sequence complementarity to the upstream fragment and reporter oligonucleotide. The template of choice can be the originally intact or glycosylase cleaved amplified nucleic acid. In addition the template can be a single stranded DNA molecule, e.g. synthetic oligonucleotide, which can vary in length and which will allow complementary annealing of the upstream fragment(s) and reporter oligonucleotide(s). The template DNA can be single stranded or double stranded in nature. Double stranded DNA acting as template consists of the template strand and the complementary strand. Double stranded DNA must first be denatured and then allowed to reanneal in the presence of the upstream fragment and reporter oligonucleotide. The upstream fragment and reporter oligonucleotide then compete with the complementary strand in annealing to the template strand of the double stranded template molecule.

Various DNA ligases and RNA ligases can be used to extend the 3' terminus of the extendible fragment by ligation of a reporter oligonucleotide on a template nucleic acid. DNA ligases from many sources, including those isolated from phage, e.g. T4 DNA ligase, and mesophilic and thermophilic bacteria, can be used to ligate the reporter oligonucleotide to the extendible fragment. Thermophilic DNA ligases can be used in repeated cyclic ligations of the extendible fragments to reporter oligonucleotides.

As discussed above, the method according to the invention involves the production of an upstream fragment which has a 3' hydroxyl group. This is an essential requirement for extension of the molecule by addition of nucleotides by a polymerase and also for extension by ligation of a reporter oligonucleotide by a ligase. In addition to a 3' hydroxyl group on the upstream fragment, the reporter oligonucleotide is required to have a 5' phosphate group for ligation to occur. Ligation, in addition, allows detection of the downstream fragment if desired. The downstream fragment is the remainder of the DNA strand from which the upstream



fragment was cleaved. Here, the reporter oligonucleotide is required to have a 3' hydroxyl terminus and the downstream fragment to have a 5' terminal phosphate group.

5            Since glycosylase mediated cleavage may generate several different  
upstream fragments in an individual reaction due to the presence of  
normal and mutant alleles in the target nucleic acid or due to the random  
introduction of modified bases, the extension by ligation reaction may  
contain several different upstream fragments in addition to several  
10 different reporter oligonucleotides and template nucleic acids. In addition  
several different nucleic acids may be characterised simultaneously since  
an individual reporter oligonucleotide and/or template nucleic acid may be  
used for the characterisation of each individual nucleic acid under  
investigation. The extended upstream fragment can be detected by any of  
15 several means including size analysis, hybridisation and amplification. In  
addition, the DNA molecule resulting from ligation can be further  
amplified in a polymerase chain reaction.

20            The reporter oligonucleotide, upstream fragment or template  
nucleic acid may be labelled. For example a fluorescent or radioactive  
label may be used in addition to a biotin or digoxigenin label. A useful  
label on the reporter oligonucleotide is a 5' terminal radioactive  
phosphate, i.e.  $^{32}\text{P}$  or  $^{33}\text{P}$ . This radioactive phosphate serves as a label with  
which to detect the DNA and also as the necessary 5' phosphate on the  
25 reporter molecule. A biotin label on the reporter oligonucleotide,  
upstream fragment or template nucleic acid will serve to immobilise the  
extended upstream fragment directly, or *via* hybridisation, to a solid  
support. Immobilisation, in combination with multiple different ligation  
extensions will serve to produce a very efficient and high throughput  
30 system for characterisation of DNA molecules.

The invention, using extension by ligation, may also be used to  
identify unknown sequence changes in nucleic acid by its ability to identify

a mutant upstream fragment in a mixture of normal upstream fragments. A partially degenerate reporter oligonucleotide may be used in the ligation reaction. Selective ligation of an upstream fragment arising from cleavage of a DNA molecule at a mutation site can also be achieved by using a  
5 degenerate reporter oligonucleotide where the 5' terminus is complementary to the normal allele. In addition the invention can be used to investigate all CpG dinucleotide sites within a DNA fragment by using a fully complementary reporter oligonucleotide designed to anneal at each CpG site. The length of any resulting ligation product will indicate the  
10 CpG site which has been mutated.

Suitably, any extended fragments resulting from step iv) are detected by hybridisation.

The method according to the invention offers significant advantages over existing methods in that

15 a) it allows detection of the similarities or the differences, or the similarities and differences between nucleic acid samples. In particular, it allows the detection of same at large numbers of multiple different loci in a nucleic acid. While it is possible to use other methods to achieve this end point, this method offers the advantage of a single process which can  
20 readily be scaled up allowing rapid and easy characterisation of nucleic acid molecules.

b) It will be appreciated that amplification of nucleic acids is a common method for characterising and detecting nucleic acids. Amplification is dependent on the use of primers which are extended in  
25 the amplification process. In addition, the method according to the invention allows the generation of high specificity primers for amplification of nucleic acids without the necessity of prior knowledge of any of the sequence of the nucleic acid. Thus, the present method has a

high utility for the characterisation of nucleic acids through analysis of amplification products generated therefrom. The amplification approaches described for characterising nucleic acids according to the invention allow characterisation of nucleic acids in a way that was not possible prior to the present invention.

c) in the field, there is a need for simplified methods for the detection of specific mutations at candidate loci. The method according to the invention offers such a simplified method for the detection of such mutations. In particular, the upstream fragment generated from sequence variation loci in nucleic acids allow the analysis of such loci using many different analytical approaches and allow accurate and simplified detection of sequence variations at such loci.

d) the method according to the invention offers a more reproducible method for characterising nucleic acids by random amplification and is less susceptible to error.

The method according to the invention can also be used to analyse the CpG content of DNA by detecting C to T transitions in DNA.

Upon analysis of mutation data and mutation spectra that have been generated over many years of mutation research, a definite hotspot for mutations in all organisms which contain 5-methylcytosine in their genomic DNA e.g. humans, has been identified, namely, mutations at CpG dinucleotides. CpG dinucleotides are a site for cytosine methylation in human cells and have been implicated in many structural and regulatory roles in genome organisation and gene expression, respectively. Cytosine in DNA is normally susceptible to a low but measurable level of deamination to form uracil, an event which is mutagenic if not repaired. However, upon methylation at the 5' site, the cytosine ring now is even more susceptible to spontaneous deamination. Therefore, the 5-

methylocytosine residue is deaminated to become thymine. Following deamination, the CpG dinucleotide becomes a TpG dinucleotide. Therefore, since 5-methylocytosine occurs only at a CpG, and the major cause of mutation at this site is due to deamination of the 5-methylocytosine, mutated CpG sequences appear in DNA as TpG dinucleotide sequences which represents a classic C to T transition mutation. Since CpG has been demonstrated to be such a hotspot for mutation in human genetic studies, a rapid screen for enhanced detection of mutations at CpG dinucleotides within the test fragment of DNA is highly advantageous.

To enhance the detection of mutations at CpG sequences, one of the following procedures can be carried out subsequent to amplification of the target DNA followed by cleavage to generate the upstream fragments.

Step iv) is carried out in the presence of a polymerase, dideoxyTTP(ddTTP), labelled dideoxy CTP(ddCTP) and wild type DNA as template, the upstream fragments are extended by incorporation of dideoxynucleotides, but only sites where C has mutated to T, including sites where CpG has mutated to TpG will become labelled following the incorporation of the labelled ddCTP. Since the ddCTP is a chain terminator nucleotide, the DNA will not be extended beyond this point. Therefore the DNA will be extended at mutated CpG sequences and will also be labelled and detectable.

Step iv) is carried out in the presence of a polymerase, wild type DNA as template, dTTP and labelled dideoxyGTP (ddGTP), the upstream fragments are extended by incorporation of dTMP, but only sites where the T is followed by G, i.e. TpG sequences will become labelled following the incorporation of the labelled ddGTP. Since the ddGTP is a chain terminator nucleotide, the DNA will not be extended beyond this point. Therefore the DNA is now extended at all TpG sequences and is also labelled and detectable. Since TpG dinucleotides occur naturally in DNA

as well as arising from CpG deamination, wild type DNA will show a characteristic banding pattern (normally only a few bands will be seen). The label can be e.g. radioactive  $^{33}\text{P}$ , fluorescent, or biotin labelled in a manner known *per se*.

- 5           Step iv) is carried out in the presence of polymerase, dUTP, dGTP, dATP, labelled dCTP and wild type DNA as template, the upstream fragments are extended by incorporation of deoxynucleotides and become labelled. The extended upstream fragments are subsequently cleaved by uracil DNA glycosylase and abasic site cleavage agent. Only that upstream  
10   fragment which was generated by cleavage at the mutation site will remain extended and labelled. This procedure will detect all mutations generating a T incorporation site including C to T mutations at CpG sites. This procedure has the advantage that step iv) utilises the incorporation of deoxynucleotides which are incorporated more efficiently than  
15   dideoxynucleotides by all polymerases.

In addition, in heterozygous samples, the mutated CpG will give rise to T/G mismatches (or U/G after introduction of dUTP as modified nucleotide). The thymine DNA glycosylase can be used to specifically cleave at these T/G or U/G mismatches at a mutated CpG sequence.

20   Brief Description of Drawings

Fig 1 is a diagram of an AP site generated during one step of the method according to the invention and of a number of ways in which the DNA at the abasic site can be cleaved so as to generate an upstream DNA fragment that can be extended;

- 25           Fig 2 is a schematic representation of the method according to the invention as described in Example 1;

Fig 3 is a schematic representation of the method according to the invention as described in Example 2 where a labelled ddTTP was used in a linear amplification reaction following generation of an upstream extendible fragment;

5            Fig 4 is a schematic representation of the method according to the invention as described in Example 2 where a labelled ddCTP was used in a linear amplification reaction following generation of an upstream extendible fragment;

10           Fig. 5 is a schematic representation of the method according to the invention as described in Example 3;

Fig. 6A-6D is a schematic representation of the extension products obtained in Example 3 following electrophoresis and autoradiography and analysis of the autoradiographs; and

15           Fig. 7A-7D is a schematic representation of the ligation reactions carried out in Example 4 on the upstream fragments and the products thereby obtained.

20           Fig. 1 depicts a single strand of DNA in 5' to 3' orientation. The position of two bases is shown, i.e. base 1 and base 3. Base 2 has been removed and an AP site exists at this position. The vertical lines denote the ribose ring which is attached to the base. The diagonal lines with 'P' in a circle refer to the phosphodiester linkages linking each ribose. The DNA can be cut at the 5' side or 3' side of an AP site as shown by arrows.

25           If the DNA is cleaved at the 3' side of the AP site, i.e. position Z, then the upstream fragment (containing base 1) has a deoxyribose phosphate moiety at its 3' terminus as shown in box C. Cleavage of the DNA at the 3' side of the AP site can be achieved by treatment with a 3' AP endonuclease / lyase or heat treatment as hereinabove described.

The AP site can also be cut in two different ways on the 5' side of the AP site, i.e. at positions X and Y. Cleavage at position X results in a OH group at the 3' terminus on the upstream fragment, as shown in box A. Cleavage at position X can be achieved by treatment with a 5' AP  
5 endonuclease as hereinabove described. Cleavage at position Y results in a phosphate group at the 3' terminus of the upstream fragment, as shown in box B. Cleavage at position Y can be achieved by treatment with heat and alkali as hereinabove described.

10 The invention will be further illustrated by the accompanying Examples.

### Modes for Carrying Out the Invention

#### Example 1

The method according to the invention was used to demonstrate the production of an extended DNA fragment by extension from the 3'OH  
15 group of an upstream fragment, which had been generated by cleavage of uracil containing DNA at the site of incorporation of the modified nucleotide. Target nucleic acid was a region of the RYR1 gene (952 to 1044) amplified from human cDNA by using upper (952 to 972) and lower (1024 to 1044) primers to generate a double stranded DNA, 93bp in  
20 length. (Nucleotide numbers refer to the sequence of the RYR1 gene)

Fig. 2 is a schematic diagram of the target nucleic acid and the upper and lower primers (primers contain standard bases, G, A, T and C). Six pmoles of the lower primer was end labelled by incubation with 1 unit of T4 Polynucleotide Kinase (commercially available from New England  
25 Biolabs), 70mM Tris-HCl (pH7.6), 10mM MgCl<sub>2</sub>, 5mM dithiothreitol and 1μCi γ<sup>32</sup>P ATP (3000Ci/mmol) for 30 min at 37°C. The target nucleic acid sample was amplified by PCR in a reaction mix containing target nucleic acid, 0.2mM dATP, dCTP, dGTP and dUTP, 6pmoles of <sup>32</sup>P labelled lower primer and non-labelled upper primer in a total volume of  
30 19μl. The reaction mix was then overlaid with an equal volume of mineral

- oil and a hot start PCR was performed whereby the reaction mix was heated to 94°C for 5min prior to addition of 1 unit of Taq polymerase (available from Promega) (bringing the total volume to 20µl). 30 cycles of denaturation, annealing and extension were carried out in a
- 5 thermocycler. The reaction mixture bearing the amplified target nucleic acid was then treated with exonuclease I (available from Amersham Life Sciences) to digest the primers not extended in the amplification step and shrimp alkaline phosphatase (SAP) (available from Boehringer Mannheim) to digest the dNTPs not incorporated during the amplification
- 10 step. This was achieved by incubating 10µl of the PCR reaction with 0.5 units of exonuclease I and 1 unit of SAP at 37°C for 30 min. Exo I and SAP were subsequently heat inactivated by incubating the reaction at 80°C for 15 min.
- 15 Uracil DNA-glycosylase (available from New England Biolabs) (0.5 units) was then added and the incubation continued at 37°C for 30 min. Following treatment with uracil DNA-glycosylase, the abasic sites generated in the amplified product were cleaved to completion by adding
- 20 NaOH to a final concentration of 0.05M and heating the mixture for 15min at 95°C. The digested DNA was then precipitated by adding 10% volume 3M sodium acetate and 2 volumes of ethanol. The pellet was resuspended in 5µl water. The digested DNA was then treated with 0.5 units of T4 polynucleotide kinase (PNK) which removes the phosphate group from the 3' terminii.
- 25 A linear amplification reaction was then carried out using the products of the above cleavage reaction, the fragment of interest being the labelled upstream extendible fragment, during which, the upstream extendible fragment is extended by a thermostable DNA polymerase in a
- 30 cycling reaction in a total volume of 10 µl. The template for this reaction



is amplified target nucleic acid (952 to 1044 of RYR1 gene) which is free from primers due to pretreatment with ExoI.

5 An equal volume of formamide loading dye (90% formamide, 0.025% bromophenol blue, 0.025% xylene cyanol) was added to the sample which was then heated at 85°C for 5 min. The sample was then loaded onto a 20% denaturing (7M urea) polyacrylamide gel and electrophoresis was carried out for 3-4 hours at 60W for size analysis of the extension products. Following electrophoresis, autoradiography was carried out by exposing the gel directly to X-ray photographic film for 12  
10 hrs at 70°C.

Analysis of the autoradiograph, where the lower primer was labelled, showed a product of 93 nucleotides in length. This product was not observed if the T4 polynucleotide kinase treatment or the linear amplification reaction was not included in the above procedure.

15 Example 2

The method according to the invention was used to detect the presence of a G to A mutation at position 1021 in the human RYR1 gene. cDNA from a normal individual and from an individual affected with  
20 Malignant Hyperthermia was amplified by using upper (952 to 972) and lower (1204 to 1224) primers to generate a 273 bp double stranded DNA fragment (952 to 1224)(As in the case of Example 1 nucleotide numbers refer to the sequence of the RYR1 gene).

25 Figs. 3 and 4 are schematic diagrams of the target nucleic acid and the upper and lower primers (primers contain standard bases, G, A, T and C). The target nucleic acid sample was amplified by PCR in a reaction mix containing target nucleic acid, 0.2mM dATP, dCTP, dGTP and 0.19mM dTTP and 0.01mM dUTP, and 6pmoles of upper and lower primers in a total volume of 19µl. The reaction mix was then overlaid with an equal

volume of mineral oil and a hot start PCR was performed whereby the reaction mix was heated to 94°C for 5min prior to addition of 1 unit of Taq polymerase (bringing the total volume to 20µl). 30 cycles of denaturation, annealing and extension were carried out in a thermocycler.

5 The reaction mixture bearing the amplified target nucleic acid was then treated with exonuclease I to digest the primers not extended in the amplification step and shrimp alkaline phosphatase (SAP) to digest the dNTPs not incorporated during the amplification step. This was achieved by incubating 10µl of the PCR reaction with 0.5 units of exonuclease I

10 and 1 unit of SAP at 37°C for 30 min. Exo I and SAP were subsequently heat inactivated by incubating the reaction at 80°C for 15 min.

Uracil DNA-glycosylase (0.5 units) was then added and the incubation continued at 37°C for 30 min. Following treatment with uracil

15 DNA-glycosylase, the abasic sites generated in the amplified product were cleaved to completion by adding NaOH to a final concentration of 0.05M and heating the mixture for 15 min at 95°C. The digested DNA was then precipitated by adding 10% volume 3M sodium acetate and 2 volumes of ethanol. The pellet was resuspended in 5µl water. The digested DNA was

20 then treated with 0.5 units of T4 Polynucleotide Kinase which removes the phosphate group from the 3' terminii.

A linear amplification reaction was then carried out using the products of the above cleavage reaction, i.e. the various extendible upstream fragments, during which, the upstream fragment is extended by

25 a thermostable DNA polymerase (i.e. Thermosequenase (available from Amersham Life Sciences)) in a cycling reaction in a total volume of 10 µl. The template for this reaction is amplified fragment of normal cDNA (952 to 1224 of RYR1 gene) which is free from unlabelled primers due to pretreatment with ExoI. The extension reaction is carried out in the

30 presence of 1mM of three of the dideoxy terminator nucleotides (ddNTP) and 0.02mM of a <sup>33</sup>P-labelled ddNTP.

An equal volume of formamide loading dye (90% formamide, 0.025% bromophenol blue, 0.025% xylene cyanol) was added to the sample which was then heated at 85°C for 5 min. The sample was then loaded onto a 6% denaturing (7M urea) polyacrylamide gel and electrophoresis was carried out for 3-4 hours at 60W for size analysis of the extension products. Following electrophoresis, autoradiography was carried out by exposing the gel directly to X-ray photographic film for 12 hrs at -70°C.

Analysis of the autoradiograph, where ddTTP was the labelled ddNTP (Fig. 3), showed a ladder of labelled fragments corresponding to the distance from the 5' end of the primers to the site of a dUMP incorporation (Only extension of the lower strand is shown). These sites corresponded to the wild type T pattern of the DNA. When mutant target nucleic acid was used to generate the extension primers, an additional band was observed in the T pattern of bands (203 nucleotides). Analysis of the autoradiograph, where ddCTP was labelled (Fig. 4) showed no labelled bands when amplified normal target nucleic acid was the source of upstream fragments, however, when amplified mutant target nucleic acid was used, analysis showed just one band. The size of the labelled fragment corresponds to the distance between the 5' end of the lower primer and the mutation site (i.e. 203 nucleotides) thereby demonstrating the presence of the G to A mutation at position 1021 of the RYR1 gene in that individual.

### Example 3

The method according to the invention was used to detect the presence of a G to A mutation at position 6411 (codon 12) in the human Ki-ras gene. Genomic DNA from normal tissue and tumour tissue of an individual with colon cancer was amplified by using upper (6390 to 6409) and lower (6417 to 6443) primers to generate a 54 bp double stranded DNA fragment (6390 to 6443)(Nucleotide numbers refer to the genomic sequence of the Ki-ras gene including introns).

Fig. 5 is a schematic diagram of the target nucleic acid and the upper and lower primers (primers contain standard bases, G, A, T and C). The target nucleic acid sample was amplified by PCR in a reaction mix containing target nucleic acid, 0.2mM dATP, dCTP, dGTP and dUTP and 6pmoles of upper and lower primers in a total volume of 19 $\mu$ l. The reaction mix was then overlaid with an equal volume of mineral oil and a hot start PCR was performed whereby the reaction mix was heated to 94°C for 5min prior to addition of 1unit of Taq polymerase (bringing the total volume to 20 $\mu$ l). 30 cycles of denaturation, annealing and extension were carried out in a thermocycler. The reaction mixture bearing the amplified target nucleic acid was then treated with Exonuclease I (Exo I) to digest the primers not extended in the amplification step and Shrimp Alkaline Phosphatase (SAP) to digest the dNTPs not incorporated during the amplification step. This was achieved by incubating 20 $\mu$ l of the PCR reaction with 0.5 units of Exo I and 1 unit of SAP at 37°C for 30 min. Exo I and SAP were subsequently heat inactivated by incubating the reaction at 80°C for 15 min. Uracil DNA-glycosylase (0.5 units) and Endonuclease IV (1unit) were then added and the incubation continued at 37°C for 30 min to allow complete excision of all the uracils present in the amplified DNA and cleavage of the resulting abasic sites to completion. This cleavage resulted in the upstream fragments having a 3' hydroxyl group at their 3' terminus which can be extended by action of a DNA polymerase.

Extension reactions were then carried out in a reaction mixture (10 $\mu$ l) containing the products of the above cleavage reaction, i.e. the extendible upstream fragments (2 $\mu$ l of cleavage reaction, approx. 1pmol of extendible fragments), and 100fmol of a synthetic template oligonucleotide, 0.2mM dATP, dTTP, and dGTP, 0.02mM dCTP, 1 $\mu$ Ci  $\alpha^{32}$ PdCTP, 6pmol reverse primer and 1unit Taq DNA polymerase. The reaction was carried out for 40 cycles of denaturation, annealing and extension.

An equal volume of formamide loading dye (90% formamide, 0.025% Bromophenol blue, 0.025% Xylene cyanol) was added to the sample which was then heated at 85°C for 5 min. The sample was then loaded onto a denaturing (7M urea) polyacrylamide gel and electrophoresis was carried out for 3-4 hours at 60W for size analysis of the extension products. Following electrophoresis, autoradiography was carried out by exposing the gel directly to X-ray photographic film for 12 hrs at -20°C.

Analysis of DNA from normal tissue results in the generation of a 37 nucleotide extendible fragment following glycosylase mediated cleavage using the above mentioned upper and lower primers (Fig. 5). Similar analysis of DNA from tumour tissue results in the generation of a 32 nucleotide extendible fragment (Fig. 5). Analysis of the autoradiograph showed a 66 nucleotide band following analysis of DNA from normal tissue and using template oligonucleotide 1 (Fig. 6A). This band was not observed when template oligonucleotide 2 was used in the above analysis (Fig. 6B). Analysis of the autoradiograph showed a 66 nucleotide band following analysis of DNA from tumour tissue and using template oligonucleotide 2 (Fig. 6C). This band was not observed when template oligonucleotide 1 was used in the above analysis (Fig. 6D). Therefore the presence of a mutation at codon 12 of the Ki-ras gene was determined by the ability of the upstream fragment to be extended on a mutant template oligonucleotide whereas the upstream fragment was not extended on a normal template oligonucleotide and *vice versa* for the absence of the mutation.

#### Example 4

Example 3 was repeated to the stage where the abasic sites had been cleaved to completion. This cleavage resulted in the upstream fragments having a 3' hydroxyl group at their 3' terminus as before.

Reporter oligonucleotides 1 and 2 (Fig. 7)(complementary to nucleotides 6397 to 6406 (5' TCCAACCTACC3' No. 1 (SEQ ID NO. 28)) and nucleotides 6402 to 6411 (5' CCAGCTCCAA3', No. 2 (SEQ ID NO. 30)) of the Ki-ras gene respectively) were 5' endlabelled using  $\gamma^{32}$ PATP and T4 polynucleotide kinase. This served to label the resulting ligated fragment and also provided a 5' terminus phosphate on the reporter oligonucleotide which is required for any possible ligation.

Ligation reactions (16°C for 60 min) were then carried out using the products of the above cleavage reaction, i.e. the extendible upstream fragments (5µl of cleavage reaction, approx. 2pmol of extendible fragments), 4pmol of labelled reporter oligonucleotide and 2pmol of a template oligonucleotide 1 or 2 (5'GGTAGTTGGAGCTGGTGGCG3' (SEQ ID No. 27) (nuc 6397 to 6416) or 5'TTGGAGCTGGTGGCGTAGGC3' (SEQ ID No. 31) (nuc 6402 to 6421) respectively) (Fig. 7) and 1 unit T4 DNA ligase, during which, the upstream fragment is extended in length by the ligation of a reporter oligonucleotide in a 20µl reaction.

An equal volume of formamide loading dye (90% formamide, 0.025% Bromophenol blue, 0.025% Xylene cyanol) was added to the sample which was then heated at 85°C for 5 min. The sample was then loaded onto a denaturing (7M urea) polyacrylamide gel and electrophoresis was carried out for 3-4 hours at 60W for size analysis of the extension products. Following electrophoresis, autoradiography was carried out by exposing the gel directly to X-ray photographic film for 3 hrs at -20°C.

Analysis of DNA from normal tissue results in the generation of a 37 nucleotide extendible fragment following glycosylase mediated cleavage using the above mentioned upper and lower primers (Fig. 5). Similar analysis of DNA from tumour tissue results in the generation of a 32 nucleotide extendible fragment (Fig. 5). Analysis of the autoradiograph

showed a 47 nucleotide band following analysis of DNA from normal tissue and using reporter oligonucleotide 1 and template oligonucleotide 1 (Fig. 7A). This band was not observed when reporter oligonucleotide 2 was used in the above analysis (Fig. 7B). Analysis of the autoradiograph  
5 showed a 42 nucleotide band following analysis of DNA from tumour tissue and using reporter oligonucleotide 2 and template oligonucleotide 2 (Fig. 7C). This band was not observed when reporter oligonucleotide 1 was used in the above analysis (Fig. 7D). Therefore the presence of a mutation at codon 12 of the Ki-ras gene was determined by the presence  
10 of a 42 nucleotide band whereas the presence of the normal allele was determined by the presence of a 47 nucleotide band. Samples containing normal and tumour DNA yielded both 42 and 47 nucleotide bands.

The above analysis was also carried out where the lower primer was  
15 5' endlabelled with  $^{32}\text{P}$ . This required that the reporter oligonucleotides be phosphorylated using unlabelled ATP as phosphate donor. The results were similar to those observed above in that a 42n band demonstrated the presence of the mutant Ki-ras gene (codon 12) whereas a 47n band demonstrated the absence of the mutation at codon 12 of the Ki-ras gene.  
20 In addition the above analysis was also carried out using the initially amplified Ki-ras gene fragment from normal or mutant sample as template during the ligation reaction. Normal amplified product was used instead of template oligonucleotide 1, whereas mutant amplified product was used instead of template oligonucleotide 2. Again the same results  
25 were observed as described above.

\*\*\*\*

As indicated above, the method according to the invention has numerous advantages over known methods, especially the method of WO  
97/03210. After the glycosylase mediated cleavage in the case of WO  
30 97/03210 (which can be carried out in a number of ways and which yields a number of different 3' termini), the resulting DNA fragments are not

further processed and are analysed directly. In the present invention, the glycosylase mediated cleavage step is followed by a step that permits extension of the 3' termini generated by glycosylase mediated cleavage.

5 A major advantage of the present invention, which is not possible in the case of the prior art methods, is that the present invention allows detection of sequence differences such as mutations and polymorphisms between nucleic acid molecules without detecting sequence similarities. As indicated above, it is not possible with the method of WO 97/03210 to  
10 detect sequence differences between nucleic acid molecules without detecting sequence similarities. This is a limitation of the method of WO 97/03210 as multiple samples cannot be combined for simultaneous analysis. The present invention also allows for the analysis of multiple genes or gene segments simultaneous analysis. Furthermore, the present invention allows for the generation of specific primers for amplification  
15 of nucleic acids without the necessity of having prior knowledge of the sequence of the nucleic acid. This is not possible with the known methods. In addition, the present invention permits the generation of specific primers from nucleic acids in a unique way and such primers can be subsequently assayed by polymerase extension to determine the nature of  
20 the sequence at the 3' termini of said primers.



## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

## (i) APPLICANT:

(A) NAME: FORBAIRT (trading as BioResearch Ireland)  
(B) STREET: Glasnevin  
(C) CITY: Dublin 9  
(E) COUNTRY: Ireland  
(F) POSTAL CODE (ZIP): none

(A) NAME: UNIVERSITY COLLEGE CORK  
(B) STREET: College Road  
(C) CITY: Cork  
(E) COUNTRY: Ireland  
(F) POSTAL CODE (ZIP): none

(A) NAME: MCCARTHY, Thomas Valentine  
(B) STREET: Vista Villa, Montenotte  
(C) CITY: Cork  
(E) COUNTRY: Ireland  
(F) POSTAL CODE (ZIP): none

(A) NAME: VAUGHAN, Patrick Martin  
(B) STREET: 175 West Avenue, Parkgate, Frankfield  
(C) CITY: Cork  
(E) COUNTRY: Ireland  
(F) POSTAL CODE (ZIP): none

(ii) TITLE OF INVENTION: A method for the characterisation of nucleic acid molecules involving generation of extendible upstream DNA fragments resulting from the cleavage of nucleic acid at an abasic site

(iii) NUMBER OF SEQUENCES: 32

## (iv) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk  
(B) COMPUTER: IBM PC compatible  
(C) OPERATING SYSTEM: PC-DOS/MS-DOS  
(D) SOFTWARE: PatentIn Release #1.0, Version #1.30 (EPO)

## (2) INFORMATION FOR SEQ ID NO: 1:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 93 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: double  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:  
(F) TISSUE TYPE: Skeletal muscle

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

TCCAAGGAGA AGCTGGATGT GGCCCCCAAG CGGGATGTGG AGGGCATGGG CCCCCCTGAG 60

ATCAAGTACG GGGAGTCACT GTGCTTCGTG CAG 93

(2) INFORMATION FOR SEQ ID NO: 2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 93 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "DNA generated by PCR amplification"

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

TCCAAGGAGA AGCTGGATGT GGCCCCCAAG CGGGAUGUGG AGGGCAUGGG CCCCCCUGAG 60

AUCAAGUACG GGGAGUCACU GUGCUUCGUG CAG 93

(2) INFORMATION FOR SEQ ID NO: 3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 93 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "DNA generated by PCR amplification"

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

CTGCACGAAG CACAGTGACT CCCCGUACUU GAUCUCAGGG GGGCCCAUGC CCUCCACAUC 60

CCGCUUGGGG GCCACAUCCA GCUUCUCCUU GGA 93

(2) INFORMATION FOR SEQ ID NO: 4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 25 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "DNA generated by glycosylase mediated cleavage and has a 3' phosphate group"

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

CTGCACGAAG CACAGTGACT CCCCG 25

(2) INFORMATION FOR SEQ ID NO: 5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 25 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "DNA generated by glycosylase mediated cleavage and has a 3' hydroxyl group"

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

CTGCACGAAG CACAGTGACT CCCCG 25

## (2) INFORMATION FOR SEQ ID NO: 6:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 93 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "DNA generated by glycosylase mediated cleavage followed by extension of upstream fragment"

## (iii) HYPOTHETICAL: NO

## (iv) ANTI-SENSE: NO

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

CTGCACGAAG CACAGTGACT CCCCGTACTT GATCTCAGGG GGGCCCATGC CCTCCACATC	60
CCGCTTGGGG GCCACATCCA GCTTCTCCTT GGA	93

## (2) INFORMATION FOR SEQ ID NO: 7:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 273 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: cDNA

## (iii) HYPOTHETICAL: NO

## (iv) ANTI-SENSE: NO

## (vi) ORIGINAL SOURCE:

(F) TISSUE TYPE: Skeletal muscle

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:

TCCAAGGAGA AGCTGGATGT GGCCCCAAG CGGGATGTGG AGGGCATGGG CCCCCCTGAG	60
ATCAAGTACG GGGAGTCACT GTGCTTCGTG CAGCATGTGG CCTCAGGACT GTGGCTCACC	120
TATGCCGCTC CAGACCCCAA GGCCCTGCGG CTCGGCGTGC TCAAGAAGAA GGCCATGCTG	180
CACCAGGAGG GCCACATGGA CGACGCACTG TCGCTGACCC GCTGCCAGCA GGAGGAGTCC	240
CAGGCCGCCC GCATGATCCA CAGCACC AAT GGC	273

## (2) INFORMATION FOR SEQ ID NO: 8:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 273 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: double  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:  
(F) TISSUE TYPE: Skeletal muscle

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:

TCCAAGGAGA AGCTGGATGT GGCCCCCAAG CGGGATGTGG AGGGCATGGG CCCCCCTGAG	60
ATCAAGTACA GGGAGTCACT GTGCTTCGTG CAGCATGTGG CCTCAGGACT GTGGCTCACC	120
TATGCCGCTC CAGACCCCAA GGCCCTGCGG CTCGGCGTGC TCAAGAAGAA GGCCATGCTG	180
CACCAGGAGG GCCACATGGA CGACGCACTG TCGCTGACCC GCTGCCAGCA GGAGGAGTCC	240
CAGGCCGCCC GCATGATCCA CAGCACCAAT GGC	273

## (2) INFORMATION FOR SEQ ID NO: 9:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 196 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid  
(A) DESCRIPTION: /desc = "DNA generated by  
glycosylase mediated cleavage and upstream fragment extension,  
and has a 3' hydrogen atom"

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(ix) FEATURE:  
(A) NAME/KEY: modified\_base  
(B) LOCATION:196  
(D) OTHER INFORMATION:/mod\_base= OTHER  
/note= "Dideoxy T"

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9:

```
GCCATTGGTG CTGTGGATCA TCGGGCGGC CTGGGACTCC TCCTGCTGGC AGCGGGTCAG      60
CGACAGTGCG TCGTCCATGT GGCCCTCCTG GTGCAGCATG GCCTTCTTCT TGAGCACGCC      120
GAGCCGCAGG GCCTTGGGGT CTGGAGCGGC ATAGGTGAGC CACAGTCCTG AGGCCACATG      180
CTGCACGAAG CACAGT                                     196
```

## (2) INFORMATION FOR SEQ ID NO: 10:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 200 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "DNA generated by glycosylase mediated cleavage followed by upstream fragment extension, and has a 3' hydrogen atom"

## (iii) HYPOTHETICAL: NO

## (ix) FEATURE:

- (A) NAME/KEY: modified\_base
- (B) LOCATION: 200
- (D) OTHER INFORMATION: /mod\_base= OTHER  
/note= "Dideoxy T"

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10:

```
GCCATTGGTG CTGTGGATCA TCGGGCGGC CTGGGACTCC TCCTGCTGGC AGCGGGTCAG      60
CGACAGTGCG TCGTCCATGT GGCCCTCCTG GTGCAGCATG GCCTTCTTCT TGAGCACGCC      120
GAGCCGCAGG GCCTTGGGGT CTGGAGCGGC ATAGGTGAGC CACAGTCCTG AGGCCACATG      180
CTGCACGAAG CACAGTGACT                                     200
```

## (2) INFORMATION FOR SEQ ID NO: 11:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 204 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid  
(A) DESCRIPTION: /desc = "DNA generated by glycosylase mediated cleavage followed by upstream fragment extension, and has a 3' hydrogen atom"

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(ix) FEATURE:

(A) NAME/KEY: modified\_base

(B) LOCATION:204

(D) OTHER INFORMATION:/note= "Dideoxy T"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 11:

GCCATTGGTG CTGTGGATCA TCGGGCGGC CTGGGACTCC TCCTGCTGGC AGCGGGTCAG	60
CGACAGTGCG TCGTCCATGT GGCCCTCCTG GTGCAGCATG GCCTTCTTCT TGAGCACGCC	120
GAGCCGCAGG GCCTTGGGGT CTGGAGCGGC ATAGGTGAGC CACAGTCCTG AGGCCACATG	180
CTGCACGAAG CACAGTGACT CCCT	204

(2) INFORMATION FOR SEQ ID NO: 12:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 206 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "DNA generated by glycosylase mediated cleavage and followed by upstream fragment extension, and has a 3' hydrogen atom"

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(ix) FEATURE:

(A) NAME/KEY: modified\_base

(B) LOCATION:206

(D) OTHER INFORMATION:/note= "Dideoxy T"

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 12:

GCCATTGGTG CTGTGGATCA TCGGGGCGGC CTGGGACTCC TCCTGCTGGC AGCGGGTCAG	60
CGACAGTGCG TCGTCCATGT GGCCCTCCTG GTGCAGCATG GCCTTCTTCT TGAGCACGCC	120
GAGCCGCAGG GCCTTGGGGT CTGGAGCGGC ATAGGTGAGC CACAGTCCTG AGGCCACATG	180
CTGCACGAAG CACAGTGA CT CCCCCT	206

## (2) INFORMATION FOR SEQ ID NO: 13:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 209 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "DNA generated by glycosylase mediated cleavage followed by upstream fragment extension, and has a 3' hydrogen atom"

## (iii) HYPOTHETICAL: NO

## (iv) ANTI-SENSE: NO

## (ix) FEATURE:

- (A) NAME/KEY: modified\_base
- (B) LOCATION: 209
- (D) OTHER INFORMATION: /mod\_base= OTHER  
/note= "Dideoxy T"

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 13:

GCCATTGGTG CTGTGGATCA TCGGGGCGGC CTGGGACTCC TCCTGCTGGC AGCGGGTCAG	60
CGACAGTGCG TCGTCCATGT GGCCCTCCTG GTGCAGCATG GCCTTCTTCT TGAGCACGCC	120
GAGCCGCAGG GCCTTGGGGT CTGGAGCGGC ATAGGTGAGC CACAGTCCTG AGGCCACATG	180
CTGCACGAAG CACAGTGA CT CCCCCTACT	209

## (2) INFORMATION FOR SEQ ID NO: 14:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 204 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear



(ii) MOLECULE TYPE: other nucleic acid  
(A) DESCRIPTION: /desc = "DNA generated by glycosylase mediated cleavage followed by upstream fragment extension, and has a 3' hydrogen atom"

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(ix) FEATURE:

(A) NAME/KEY: modified\_base

(B) LOCATION:204

(D) OTHER INFORMATION:/mod\_base= OTHER  
/note= "Dideoxy C"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 14:

GCCATTGGTG CTGTGGATCA TGCGGGCGGC CTGGGACTCC TCCTGCTGGC AGCGGGTCAG	60
CGACAGTGC GCGTCCATGT GGCCCTCCTG GTGCAGCATG GCCTTCTTCT TGAGCACGCC	120
GAGCCGCAGG GCCTTGGGGT CTGGAGCGGC ATAGGTGAGC CACAGTCCTG AGGCCACATG	180
CTGCACGAAG CACAGTGA CTCCCC	204

(2) INFORMATION FOR SEQ ID NO: 15:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 54 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: double

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 15:

AACTTGTGGT AGTTGGAGCT GGTGGCCTAG GCAAGAGTGC CTTGACGATA CAGC	54
---	----

## (2) INFORMATION FOR SEQ ID NO: 16:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 54 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "Generated by PCR  
amplification of genomic DNA"

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 16:

AACTTGTTGGT AGTTGGAGCT GGUGGCCGUAG GCAAGAGUGC CUUGACGAUA CAGC

54

## (2) INFORMATION FOR SEQ ID NO: 17:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 54 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "Generated by PCR  
amplification of genomic DNA"

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 17:

GCTGTATCGT CAAGGCACTC TTGCCTACGC CACCAGCUCC AACUACCACA AGUU

54

## (2) INFORMATION FOR SEQ ID NO: 18:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 54 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "Generated by PCR  
amplification of genomic DNA"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 18:

AACTTGTGGT AGTTGGAGCT GAUGGCGUAG GCAAGAGUGC CUUGACGAUA CAGC

54

(2) INFORMATION FOR SEQ ID NO: 19:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 54 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "Generated by PCR amplification"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 19:

GCTGTATCGT CAAGGCACTC TTGCCTACGC CAUCAGCUCC AACUACCACA AGUU

54

(2) INFORMATION FOR SEQ ID NO: 20:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 37 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "Generated by glycosylase mediated cleavage of PCR amplified DNA"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 20:

GCTGTATCGT CAAGGCACTC TTGCCTACGC CACCAGC

37

(2) INFORMATION FOR SEQ ID NO: 21:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 32 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid  
(A) DESCRIPTION: /desc = "Generated by glycosylase  
mediated cleavage of PCR amplified DNA"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 21:

GCTGTATCGT CAAGGCACTC TTGCCTACGC CA

32

(2) INFORMATION FOR SEQ ID NO: 22:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 66 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid  
(A) DESCRIPTION: /desc = "synthetic oligonucleotide"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 22:

GCTGTAAACG ACGGCCAGTT TCATGCAGGG CTGGAGTCGT AGGCAAGAGT GCCTTGACGA  
TACAGC

60

66

(2) INFORMATION FOR SEQ ID NO: 23:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 24 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid  
(A) DESCRIPTION: /desc = "synthetic oligonucleotide"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 23:

GCTGTAAACG ACGGCCAGTT TCAT

24

## (2) INFORMATION FOR SEQ ID NO: 24:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 66 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: other nucleic acid

- (A) DESCRIPTION: /desc = "Generated by primer extension"

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 24:

GCTGTATCGT CAAGGCACTC TTGCCTACGC CACCAGCCCT GCATGAAACT GGCCGTCGTT 60  
TACAGC 66

## (2) INFORMATION FOR SEQ ID NO: 25:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 66 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: other nucleic acid

- (A) DESCRIPTION: /desc = "synthetic oligonucleotide"

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 25:

GCTGTAAACG ACGGCCAGTT TCATGCAGGA TCCATGGCGT AGGCAAGAGT GCCTTGACGA 60  
TACAGC 66

## (2) INFORMATION FOR SEQ ID NO: 26:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 66 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: other nucleic acid

- (A) DESCRIPTION: /desc = "Generated by primer extension"

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 26:

GCTGTATCGT CAAGGCACTC TTGCCTACGC CATGGATCCT GCATGAAACT GGCCGTCGTT 60  
TACAGC 66

## (2) INFORMATION FOR SEQ ID NO: 27:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: other nucleic acid

- (A) DESCRIPTION: /desc = "synthetic oligonucleotide"

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 27:

GGTAGTTGGA GCTGGTGGCG 20

## (2) INFORMATION FOR SEQ ID NO: 28:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 10 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: other nucleic acid

- (A) DESCRIPTION: /desc = "synthetic oligonucleotide"

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 28:

TCCAACTACC 10

## (2) INFORMATION FOR SEQ ID NO: 29:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 47 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid  
(A) DESCRIPTION: /desc = "Generated by ligation of  
two DNA molecules"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 29:

GCTGTATCGT CAAGGCACTC TTGCCTACGC CACCAGCTCC AACTACC

47

(2) INFORMATION FOR SEQ ID NO: 30:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 10 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid  
(A) DESCRIPTION: /desc = "synthetic oligonucleotide"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 30:

CCAGCTCCAA

10

(2) INFORMATION FOR SEQ ID NO: 31:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 20 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid  
(A) DESCRIPTION: /desc = "synthetic oligonucleotide"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 31:

TTGGAGCTGG TGGCGTAGGC

20

(2) INFORMATION FOR SEQ ID NO: 32:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 42 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid  
(A) DESCRIPTION: /desc = "Generated by ligation of  
two DNA molecules"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 32:

GCTGTATCGT CAAGGCACTC TTGCCTACGC CACCAGCTCC AA

42



Claims:-

1. A method for characterising nucleic acid molecules, which comprises the steps of:
  - 5 i) introducing a modified base which is a substrate for a DNA glycosylase into a DNA molecule;
  - ii) excising the modified base by means of said DNA glycosylase so as to generate an abasic site;
  - iii) cleaving the DNA at the abasic site so as to generate an upstream DNA fragment that can be extended; and
  - 10 iv) incubating the extendible upstream fragment in the presence of an enzyme allowing for extension thereof and a template nucleic acid and analysing the resultant fragment(s).
2. A method according to Claim 1, wherein the upstream  
15 fragment is generated by cleaving the DNA at the 5' side of the abasic site, such that the 3' terminus of the upstream fragment bears a hydroxyl group.
3. A method according to Claim 2, wherein the cleavage is achieved with a 5' AP endonuclease.
- 20 4. A method according to Claim 1, wherein the upstream fragment is generated by cleaving at the 5' side of the abasic site so as to leave a phosphate group at the 3' terminus of the upstream fragment and removing the phosphate group so that the upstream fragment bears a hydroxyl group at the 3' terminus.

5. A method according to Claim 1, wherein the upstream fragment is generated by cleaving at the 3' side of the abasic site so as to generate a deoxyribose phosphate group at the 3' terminus of the upstream fragment and subsequently removing the deoxyribose group to leave a hydroxyl group at the 3' terminus.
6. A method according to any preceding claim, wherein 5' deoxyribose moieties downstream of the 3' terminus of the upstream fragment are removed so that the upstream fragment can be extended on the template.
7. A method according to Claim 6, wherein the 5' deoxyribose moieties are removed by a 5' deoxyribophosphodiesterase.
8. A method according to any preceding claim, wherein the modified base is introduced by enzymatic amplification of the DNA.
9. A method according to Claim 8, wherein the amplified strands are separated for a separate analysis of the respective strands.
10. A method according to Claim 8 or 9, wherein a primer or one or more nucleotide(s) involved in the enzymatic amplification is labelled.
11. A method according to any preceding claim, wherein the enzyme is a polymerase.
12. A method according to Claim 11, wherein the extendible upstream fragment is incubated with the polymerase in the presence of one or more nucleotide(s).
13. A method according to Claim 12, wherein one or more of the nucleotide(s) of step iv) is a dideoxy nucleotide.

14. A method according to Claim 12 or 13, wherein one or more of the nucleotide(s) of step iv) is labelled.

15. A method according to any one of Claims 11-14, wherein the extension of step iv) is achieved by means of an amplification reaction using said extendible DNA fragment.

16. A method according to any one of Claims 11-15, wherein the extension of step iv) is achieved by means of an amplification reaction including a primer in addition to using said extendible DNA fragment.

17. A method according to any one of Claims 1-10, wherein the enzyme is a ligase.

18. A method according to Claim 17, wherein the extendible upstream fragment is incubated with the ligase in the presence of a reporter oligonucleotide.

19. A method according to Claim 18, wherein the reporter oligonucleotide is partially degenerate.

20. A method according to any preceding claim, wherein any extended fragments resulting from step iv) are detected by hybridisation.

21. A method according to any preceding claim, which is used to detect a known or unknown mutation.

22. A method according to Claim 1 for characterising nucleic acid molecules, substantially as hereinbefore described and exemplified.

23. A method according to any one of Claims 1-20, wherein the method is used to analyse the CpG content of DNA by detecting C to T transitions in DNA.

24. A method according to Claim 1 for characterising nucleic acid molecules, substantially as hereinbefore described and exemplified.

1/7







FIG 1

2/7





FIG 2


3/7

- Amplification of normal and mutant target nucleic acid in the presence of dGTP, dATP, dCTP and 1/20 ratio of dUTP to dTTP.



- Treatment with Exonuclease I and Shrimp Alkaline Phosphatase.
- Treatment with UDG.
- Treatment with NaOH at 95°C
- DNA is precipitated.
- Treatment with T4 PNK

Extension of the upstream fragments generated above in the presence of  $^{32}\text{P}$ -labelled ddTTP\* and unlabelled ddGTP, ddATP and ddCTP.



Only some fragments corresponding to cleavage at U incorporation sites surrounding the mutation site are shown here.

Detection of extended labelled fragments by PAGE and autoradiography





FIG 3


4/7

- Amplification of normal and mutant target nucleic acid in the presence of dGTP, dATP, dCTP and 1/20 ratio of dUTP to dTTP.



- Treatment with Exonuclease I and Shrimp Alkaline Phosphatase.
- Treatment with UDG.
- Treatment with NaOH at 95°C
- DNA is precipitated.
- Treatment with T4 PNK

Extension of the upstream fragments generated above in the presence of <sup>32</sup>P-labelled ddCTP\* and unlabelled ddGTP, ddATP and ddTTP.



Only some fragments corresponding to cleavage at U incorporation sites surrounding the mutation site are shown here.



Detection of extended labelled fragments by PAGE and autoradiography

FIG 4



5/7

6390                      Upper primer                      Mutation site                      SEQ ID NO:15                      6443

5' AACTTGTGGTAGTTGGAGCTGGTGGCGTAGGCAAGAGTGCCTTGACGATACAGC 3'

3' TTGAACACCATCAACCTCGACCAACCGCATCCGTTCTCAGGAACTGCTATGTCG 5'

T                      Lower primer



Amplification of target nucleic acid in the presence of dGTP, dATP, dUTP and dCTP.

SEQ ID NO:16                      Amplified normal allele

5' AACTTGTGGTAGTTGGAGCTGGUGGCGUAGGCAAGAGUGCCUUGACGAUACAGC 3'

3' UUGAACACCAUCAACCUCGACCAACCGCATCCGTTCTCAGGAACTGCTATGTCG 5'

SEQ ID NO:17

SEQ ID NO:18                      Amplified mutant allele

5' AACTTGTGGTAGTTGGAGCTGAUGGCGUAGGCAAGAGUGCCUUGACGAUACAGC 3'

3' UUGAACACCAUCAACCUCGACUACCGCATCCGTTCTCAGGAACTGCTATGTCG 5'

SEQ ID NO:19



- Treatment with Exonuclease I and Shrimp Alkaline Phosphatase.  
- Treatment with UDG.  
- Treatment with Endo IV.

SEQ ID NO:20                      Normal upstream fragment

3' CGACCACCGCATCCGTTCTCAGGAACTGCTATGTCG 5'

SEQ ID NO:21                      Mutant upstream fragment

3' ACCGCATCCGTTCTCAGGAACTGCTATGTCG 5'

FIG 5

6/7




FIG 6

7/7




FIG 7

# INTERNATIONAL SEARCH REPORT

Int. Appl. No.

PCT/IE 98/00030

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 6 C12Q1/68

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
IPC 6 C12Q

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	VAUGHAN P ET AL: "A novel process for mutation detection using uracil DNA-glycosylase" NUCLEIC ACIDS RESEARCH, vol. 26, no. 12, February 1998, pages 810-15, XP002088826 see the whole document	1-16, 20-24
Y	WO 97 03210 A (FORFAS TRADING AS BIORESEARCH ;UNIV CORK (IE); MCCARTHY THOMAS VAL) 30 January 1997 cited in the application see the whole document	1, 17-19
Y	WO 97 12061 A (EPICENTRE TECHNOLOGIES CORP) 3 April 1997 see the whole document	1-24
Y	WO 97 12061 A (EPICENTRE TECHNOLOGIES CORP) 3 April 1997 see the whole document	1-24

-/--

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

23 December 1998

Date of making of the international search report

15/01/1999

Name and mailing address of the ISA  
European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040. Tx. 31 651 epo nl.  
Fax: (+31-70) 340-3016

Authorized officer

Osborne, H

# INTERNATIONAL SEARCH REPORT

Inter: nat Application No

PCT/IE 98/00030

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>MCGRATH A ET AL: "A DNA glycolase-based fingerprint for accurate identification of amplified DNA products and its application in the accurate diagnosis of infectious organisms"</p> <p>ANALYTICAL BIOCHEMISTRY, vol. 259, no. 2, - June 1998 pages 288-92, XP002088827 see the whole document</p>	1-24
A	<p>WO 96 30545 A (APPLIED GENETICS INC) 3 October 1996</p>	

1

# INTERNATIONAL SEARCH REPORT

Information on patent family members

Intern. Patent Application No

PCT/IE 98/00030

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9703210 A	30-01-1997	AU 4399596 A CA 2226542 A EP 0843735 A NO 980105 A	10-02-1997 30-01-1997 27-05-1998 10-03-1998
WO 9712061 A	03-04-1997	AU 7118396 A CA 2233079 A EP 0854936 A	17-04-1997 03-04-1997 29-07-1998
WO 9630545 A	03-10-1996	AU 5296496 A	16-10-1996